arXiv:1403.0299v1 [math.DG] 3 Mar 2014

ON THE FUNCTIONAL BLASCHKE-SANTALÓ INEQUALITY

YOUJIANG LIN AND GANGSONG LENG

Abstract. In this paper, using functional Steiner symmetrizations, we show that Meyer and Pajor's proof of the Blaschke-Santaló inequality can be extended to the functional setting.

1. Introduction

For a convex body $K \subset \mathbb{R}^n$ and a point $z \in \mathbb{R}^n$, the polar body K^z of K with respect to z is the convex set defined by $K^z = \{y \in \mathbb{R}^n : \langle y-z, x-z\rangle \leq 1 \text{ for every } x \in K\}$. The *Santaló point* $s(K)$ of K is a point for which $V_n(K^{s(K)}) = \min_{z\in int(K)} V_n(K^z)$, where $V_n(K)$ denotes the volume of set K. The Blaschke-Santaló inequality [4, 18, 19] states that $V_n(K)V_n(K^{s(K)}) \leq V_n(B_2^n)^2$, where B_2^n is the Euclidean ball.

For a log-concave function $f : \mathbb{R}^n \to [0,\infty)$ and a point $z \in \mathbb{R}^n$, its polar with respect to z is defined by $f^z(y) = \inf_{x\in\mathbb{R}^n} \frac{e^{-\langle x-z, y-z\rangle}}{f(x)}$. The *Santaló point* $s(f)$ of f is the point z_0 satisfying $\int f^{z_0} = \inf_{z\in\mathbb{R}^n} \int f^z$.

The *functional Blaschke-Santaló inequality* of log-concave functions is the analogue of Blaschke-Santaló inequality of convex bodies.

Theorem 1.1. *(Artstein, Klartag, Milman). Let $f : \mathbb{R}^n \to [0,+\infty)$ be a log-concave function such that $0 < \int f < \infty$. Then, $\int_{\mathbb{R}^n} f \int_{\mathbb{R}^n} f^{s(f)} \leq (2\pi)^n$ with equality holds exactly for Gaussians.*

Key words and phrases. Convex body; Polar body; Parallel sections homothety bodies; Mahler conjecture; Cylinder.

2010 Mathematics Subject Classification. 52A10, 52A40.

The authors would like to acknowledge the support from China Postdoctoral Science Foundation Grant 2013M540806, National Natural Science Foundation of China under grant 11271244 and National Natural Science Foundation of China under grant 11271282 and the 973 Program 2013CB834201.

When f is even, the functional Blaschke-Santaló inequality follows from an earlier inequality of Ball [2]; and in [9], Fradelizi and Meyer proved something more general (see also [11]). Lutwak and Zhang [13] and Lutwak et al. [14] gave other very different forms of the Blaschke-Santaló inequality. In this paper, we give a more general result than Theorem 1.1, which becomes into a special case of $\lambda = 1/2$ in Theorem 1.2.

Theorem 1.2. *Let $f : \mathbb{R}^n \to [0, +\infty)$ be a log-concave function such that $0 < \int f < \infty$. Let H be an affine hyperplane and let H_+ and H_- denote two closed half-spaces bounded by H. If $\lambda \in (0, 1)$ satisfies $\lambda \int_{\mathbb{R}^n} f = \int_{H_+} f$. Then there exists $z \in H$ such that*

$$\int_{\mathbb{R}^n} f \int_{\mathbb{R}^n} f^z \leq \frac{1}{4\lambda(1-\lambda)}(2\pi)^n. \tag{1.1}$$

In [12], Lehec proved a very general functional version for non-negative Borel functions, Theorem 1.2 is a particular case of result of Lehec. Lehec's proof is by induction on the dimension, and the proof is by functional Steiner symmetrizations. In fact, Mayer and Pajor [15] have proved the Blaschke-Santaló inequality for convex bodies, here we show that Meyer and Pajor's proof of the Blaschke-Santaló inequality can be extended to the functional setting. It has recently come to our attention that in a remark of [9], Fradelizi and Meyer expressed the same idea to prove the functional Blaschke-Santaló inequality.

2. Notations and background materials

Let $|\cdot|$ denote the Euclidean norm. Let $\mathrm{int}A$ denote the interior of $A \subset \mathbb{R}^n$. Let $\mathrm{cl}A$ denote the closure of A. Let $\dim A$ denote the dimension of A. A set $C \subset \mathbb{R}^n$ is called a *convex cone* if C is convex and nonempty and if $x \in C$, $\lambda \geq 0$ implies $\lambda x \in C$. We define $C^* := \{x \in \mathbb{R}^n : \langle x, y\rangle \leq 0 \quad \text{for all } y \in C\}$ and call this the *dual cone* of C.

For a non-empty convex set $K \subset \mathbb{R}^n$ and an affine hyperplane H with unit normal vector u, the *Steiner symmetrization* $S_H K$ of K

with respect to H is defined as $S_H K := \{x' + \frac{1}{2}(t_1 - t_2)u : \ x' \in P_H(K), \ t_i \in I_K(x') \text{ for } i = 1, 2\}$, where $P_H(K) := \{x' \in H : x' + tu \in K \text{ for some } t \in \mathbb{R}\}$ is the projection of K onto H and $I_K(x') := \{t \in \mathbb{R} : \ x' + tu \in K\}$.

Let $\bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$. For a given function $f : \mathbb{R}^n \to \bar{\mathbb{R}}$ and for $\alpha \in \bar{\mathbb{R}}$ we use the abbreviation $\{f = \alpha\} := \{x \in \mathbb{R}^n : f(x) = \alpha\}$, and $\{f \leq \alpha\}$, $\{f < \alpha\}$ etc. are defined similarly. A function $f : \mathbb{R}^n \to \bar{\mathbb{R}}$ is called *proper* if $\{f = -\infty\} = \emptyset$ and $\{f = \infty\} \neq \mathbb{R}^n$. A function ϕ is called *convex* if ϕ is proper and $\phi(\alpha x + (1-\alpha)y) \leq \alpha\phi(x) + (1-\alpha)\phi(y)$ for all $x, y \in \mathbb{R}^n$ and for any $0 \leq \lambda \leq 1$. A function f is called *log-concave* if $f = e^{-\phi}$, where ϕ is a convex function. A function $f : \mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ is called *coercive* if $\lim_{|x| \to +\infty} f(x) = +\infty$. A function f is called *symmetric* about H if for any $x' \in H$ and $t \in \mathbb{R}$, $f(x' + tu) = f(x' - tu)$. A function $f : \mathbb{R}^n \to \mathbb{R}$ is called *unconditional* about z if $f(x_1 - z_1, \ldots, x_n - z_n) = f(|x_1 - z_1|, \ldots, |x_n - z_n|)$ for every $(x_1, \ldots, x_n) \in \mathbb{R}^n$. If $z = 0$, then f is called *unconditional*.

The *effective domain* of convex function ϕ is the nonempty set $\mathrm{dom}\phi := \{\phi < \infty\}$. The *support* of function f is the set $\mathrm{supp} f := \{f \neq 0\}$. For log-concave function $f = e^{-\phi}$, it is clear that $\mathrm{supp} f = \mathrm{dom}\phi$. The nonempty set $\mathrm{epi}\phi := \{(x, r) \in \mathbb{R}^n \times \mathbb{R} : \ r \geq \phi(x)\}$ denote the *epigraph* of convex function ϕ.

For an affine subspace G of $\mathbb{R}^n$, let $G^\perp$ denote the *orthogonal complement* of G, we have $G^\perp = \{x \in \mathbb{R}^n : \langle x, y - y' \rangle = 0 \text{ for every } y, y' \in G\}$. The *Santaló point* $s_G(f)$ of f about G is a point satisfying $\int f^{s_G(f)} = \inf_{z \in G} \int f^z$. Let f be a log-concave function such that $0 < \int f < \infty$, and let H_+ and H_- be two half-spaces bounded by an affine hyperplane H; let $0 < \lambda < 1$; we shall say that H is λ-*separating* for f if $\int_{H_+} f \int_{H_-} f = \lambda(1 - \lambda) \left(\int_{\mathbb{R}^n} f\right)^2$ and when $\lambda = 1/2$, we shall say that H is *medial* for f. For a function $\phi : \mathbb{R}^n \to \bar{\mathbb{R}}$, its Legendre transform about z is defined by $\mathcal{L}^z \phi(y) = \sup_{x \in \mathbb{R}^n} [\langle x - z, y - z \rangle - \phi(x)]$. If $f(x) = e^{-\phi(x)}$, where $\phi(x)$ is a convex function, then $f^z(y) = e^{-\mathcal{L}^z \phi(y)}$.

Since $\mathcal{L}^z(\mathcal{L}^z\phi) = \phi$ for a convex function ϕ, $(f^z)^z = f$. If $z = 0$, we shall use the simpler notation $\mathcal{L}$ for $\mathcal{L}^0$.

Given two functions $f, g : \mathbb{R}^n \to [0, \infty)$, their *Asplund product* is defined by $(f \star g)(x) = \sup_{x_1+x_2=x} f(x_1)g(x_2)$. The *$\lambda$-homothety* of a function f is defined as $(\lambda \cdot f)(x) = f^\lambda(\frac{x}{\lambda})$. Then, the classical Prékopa inequality (see Prékopa [16, 17]) can be stated as follows: Given $f, g : \mathbb{R}^n \to [0, +\infty)$ and $0 < \lambda < 1$, $\int(\lambda \cdot f) \star ((1-\lambda) \cdot g) \geq \left(\int f\right)^\lambda \left(\int g\right)^{1-\lambda}$. The following lemma, as a particular case of a result due to Ball [3], was proved by Meyer and Pajor in [15].

Lemma 2.1. [15] *Let f_0, f_1, $f_2 : \mathbb{R}^+ \to \mathbb{R}^+$ be three functions such that $0 < \int_0^{+\infty} f_i < \infty$, $i = 0, 1, 2$, they are continuous and suppose that $f_0\left(\frac{2xy}{x+y}\right) \geq f_1(x)^{\frac{y}{x+y}} f_2(y)^{\frac{x}{x+y}}$ for every $x, y > 0$. Then one has*

$$\frac{1}{\int_0^{+\infty} f_0(t)dt} \leq \frac{1}{2}\left(\frac{1}{\int_0^{+\infty} f_1(t)dt} + \frac{1}{\int_0^{+\infty} f_2(t)dt}\right).$$

3. The functional Steiner symmetrization

The familiar definition of Steiner symmetrization for a nonnegative measurable function f can be stated as following (see [5, 6, 7, 8]):

Definition 1. For a measurable function $f : \mathbb{R}^n \to [0, +\infty)$ and an affine hyperplane $H \subset \mathbb{R}^n$, let m denote the Lebesgue measure, if $m(\{f > t\}) < +\infty$ for all $t > 0$, then its Steiner symmetrization is defined as

$$S_H f(x) = \int_0^\infty \mathcal{X}_{S_H\{f>t\}}(x)dt, \tag{3.1}$$

where $\mathcal{X}_A$ denotes the characteristic function of set A.

Next, we give a approach of defining Steiner symmetrization for coercive convex functions by the Steiner symmetrization of epigraphs. A similar functional steiner symmetrization is defined in a remark of AKM's paper [1] and studied in an article by Lehec [10]. The idea of our definition is same as the given definition in a remark at the end of an article by Fradelizi and Meyer [9].

Definition 2. For a coercive convex function ϕ and an affine hyperplane $H \subset \mathbb{R}^n$, we define the *Steiner symmetrization* $S_H\phi$ of ϕ with respect to H as a function satisfying

$$\mathrm{epi}(S_H\phi) = S_{\widetilde{H}}(\mathrm{cl}\ \mathrm{epi}\phi), \tag{3.2}$$

where $\widetilde{H} = \{(x', s) \in \mathbb{R}^{n+1} : x' \in H\}$ is an affine hyperplane in $\mathbb{R}^{n+1}$.

Remark 1. *(i) By Definition 2, for an integrable log-concave function $f = e^{-\phi}$, the Steiner symmetrization of f can be defined as $S_H f := e^{-(S_H\phi)}$. If we define $S_H f$ by Definition 1, then $S_H f$ still satisfies (3.2). Thus, for integrable log-concave functions, the two definitions are essentially same.*

(ii) By Definition 2, for a given $x' \in H$ and any $s \in \mathbb{R}$, we have $V_1(\{(S_H\phi)(x' + tu) < s\}) = V_1(\{\phi(x' + tu) < s\})$. By the Fubini's theorem, we have

$$\int_{\mathbb{R}} (S_H f)(x' + tu)dt = \int_{\mathbb{R}} f(x' + tu)dt. \tag{3.3}$$

Similarly, $\int_{\mathbb{R}^n} S_H f = \int_{\mathbb{R}^n} f$ is also established.

Proposition 1. For a coercive convex function ϕ and an affine hyperplane $H \subset \mathbb{R}^n$ with outer unit normal vector u, then $S_H\phi$ has the following properties.

(i) $S_H\phi$ is a closed coercive convex function and symmetric about H.

(ii) Let H_1 and H_2 be two orthogonal hyperplanes in $\mathbb{R}^n$, then $S_{H_2}(S_{H_1}\phi)$ is symmetric about both H_1 and H_2.

(iii) For any given $x' \in H$ and $t \in \mathbb{R}$, let $\phi_1(t) := \phi(x' + tu)$ and $(S\phi_1)(t) := (S_H\phi)(x' + tu)$, then $(S\phi_1)(t)$ satisfies one of the following three cases. *1).* $(S\phi_1)(t) = \phi_1(t_1) = \phi_1(t_1 - 2t)$ for some $t_1 \in \mathbb{R}$. *2).* $(S\phi_1)(t) = \phi_1(t_0 - 2t) \geq \lim_{t\to t_0,\, t<t_0} \phi_1(t)$ for some $t_0 \in \mathbb{R}$. *3).* $(S\phi_1)(t) = \phi_1(t_0 + 2t) \geq \lim_{t\to t_0,\, t>t_0} \phi_1(t)$ for some $t_0 \in \mathbb{R}$.

Proof. (i) By the fact that ϕ is convex if and only if epiϕ is convex, since ϕ is convex, epiϕ is a convex subset of $\mathbb{R}^{n+1}$. Since the closure of a convex set is convex, and the Steiner symmetrization of a convex set is

also convex, by (3.2), epi$(S_H\phi)$ is a convex subset of $\mathbb{R}^{n+1}$. Therefore, $S_H\phi$ is a convex function. By Definition 2, it is clear that $S_H\phi$ is closed, coercive and symmetric with respect to H.

(ii) Since epi$(S_{H_2}(S_{H_1}\phi))$ is symmetric about both $\widetilde{H_1}$ and $\widetilde{H_2}$, where $\widetilde{H_i} = \{(x', s) \in \mathbb{R}^{n+1} : x' \in H_i\}$ $(i = 1, 2)$, $S_{H_2}(S_{H_1}\phi)$ is symmetric about both H_1 and H_2.

(iii) If dom$\phi_1 = \mathbb{R}$, by (3.2) in Definition 2, we have

$$\text{epi}(S\phi_1) = S_{\widetilde{H}}(\text{cl epi}\phi_1). \tag{3.4}$$

Thus there exists some $t_1 \in \mathbb{R}$ satisfying

$$(S\phi_1)(t) = \phi_1(t_1) = \phi_1(t_1 - 2t). \tag{3.5}$$

If dom$\phi_1 \neq \mathbb{R}$, then there exist eight cases for domϕ_1: 1) $[\alpha, \beta]$; 2) (α, β); 3) $(\alpha, \beta]$; 4) $[\alpha, \beta)$; 5) $(-\infty, \beta]$; 6) $(-\infty, \beta)$; 7) $[\alpha, +\infty)$; 8) $(\alpha, +\infty)$. Here, we only prove our conclusion for dom$\phi_1 = (\alpha, \beta)$. By the same method we can prove our conclusion for other cases. For dom$\phi_1 = (\alpha, \beta)$, by Definition 2, it is clear that $(S\phi_1)(t) = +\infty$ for $|t| \geq \frac{\beta-\alpha}{2}$. If $|t| < \frac{\beta-\alpha}{2}$, let $\lim_{x\to\alpha,\, x>\alpha} \phi_1(x) = b_1$, $\lim_{x\to\beta,\, x<\beta} \phi_1(x) = b_2$, then we consider the following four cases. (a) If $b_1 = b_2 = +\infty$, then by (3.4), there exists some $t_1 \in \mathbb{R}$ satisfying (3.5). (b) If $b_1 < +\infty$, $b_2 = +\infty$, then there exists $\gamma \in (\alpha, \beta)$ such that $\phi_1(\gamma) = b_1$. Then by (3.4), for $|t| < \frac{\gamma-\alpha}{2}$, (3.5) is established, for $|t| \geq \frac{\gamma-\alpha}{2}$, we have $(S\phi_1)(t) = \phi_1(\alpha + 2t) \geq b_1$. (c) If $b_1 = +\infty$, $b_2 < +\infty$, then there exists $\gamma \in (\alpha, \beta)$ such that $\phi_1(\gamma) = b_2$. Then by (3.4), for $|t| < \frac{\beta-\gamma}{2}$, (3.5) is established, for $|t| \geq \frac{\gamma-\alpha}{2}$, we have $(S\phi_1)(t) = \phi_1(\beta - 2t) \geq b_2$. (d) If $b_1 < \infty$, $b_2 < +\infty$, we consider three cases. If $b_1 = b_2$, then (3.5) is established. If $b_1 > b_2$, the proof is same as in (c). If $b_1 < b_2$, the proof is same as in (b). This completes the proof. □

4. The proofs of theorems

In order to prove theorems stated in the introduction, we have to establish the following six lemmas:

Lemma 4.1. *If f be a log-concave function such that $0 < \int f < \infty$, then the function F defined by $F(z) := \int_{\mathbb{R}^n} f^z(x)dx$ has the following properties. (i) $F(z)$ is a coercive convex function on $\mathbb{R}^n$ and is strictly convex on* int dom$F$*; (ii) If $f(x)$ is even about z_0, then $F(z)$ is also even about z_0.*

Proof. (i) **Step 1.** We shall prove F is coercive. Let $f = e^{-\phi}$, for any given $z \in \mathbb{R}^n$ and $r > 0$, we have

$$F(z) = \int_{\mathbb{R}^n} f^z(x+z)dx \geq \int_{rB_2^n} f^z(x+z)dx = \int_{rB_2^n} e^{-\mathcal{L}\phi(x)+\langle x,z\rangle}dx \quad (4.1)$$

Since $f = e^{-\phi}$ is integrable, there is $\gamma > 0$ and $h \in \mathbb{R}$ such that

$$\phi(x) \geq \gamma \sum_{i=1}^{n} |x_i| + h \ \text{ for any } x \in \mathbb{R}^n. \quad (4.2)$$

Thus, for $y \in \gamma B_\infty^n$, where $B_\infty^n = \{x \in \mathbb{R}^n : |x_i| \leq 1, i = 1, \ldots, n\}$, $\mathcal{L}\phi(y) \leq \sup_{x\in\mathbb{R}^n}[\langle y, x\rangle - \gamma \sum_{i=1}^n |x_i| - h] \leq -h$. Let $rB_2^n \subset \frac{1}{2}\gamma B_\infty^n$, we have $rB_2^n \subset \text{int}(\text{dom}\mathcal{L}\phi)$. Since function $g(x) := e^{-\mathcal{L}\phi(x)}$ is continuous on rB_2^n. Thus, there exists $m > 0$ such that $g(x) \geq m$ for any $x \in rB_2^n$. Therefore,

$$\int_{rB_2^n} e^{-\mathcal{L}\phi(x)+\langle x,z\rangle}dx \geq m \int_{rB_2^n} e^{\langle x,z\rangle}dx. \quad (4.3)$$

For any $z \in \mathbb{R}^n$ and $|z| \geq 1$, let $z' = \frac{r}{2}\frac{z}{|z|}$, we get a closed half-space $H^+ = \{x \in \mathbb{R}^n : \langle x - z', z\rangle \geq 0\}$. For any $x \in H^+$, we have $\langle x, z\rangle \geq \langle z', z\rangle = \frac{r}{2}|z|$. Therefore,

$$\int_{rB_2^n} e^{\langle x,z\rangle}dx \ \geq \ \int_{(rB_2^n)\cap H^+} e^{\frac{r|z|}{2}}dx = V_n((rB_2^n)\cap H^+)e^{\frac{r|z|}{2}}. \quad (4.4)$$

Since $V_n((rB_2^n)\cap H^+)$ is a positive constant independent of z, by (4.1), (4.3) and (4.4), $F(z)$ is coercive.

Step 2. We shall prove that F is convex and is strictly convex on int dom$F$. First, we prove $F(z)$ is proper. It is clear that $F(z) > -\infty$ for any $z \in \mathbb{R}^n$. The following claim shows that $\{F = \infty\} \neq \mathbb{R}^n$.

Claim 1. For any $z \in$ int supp$f$, $F(z) < \infty$.

Proof of Claim 1. For any $z \in \text{int supp}f$, there is a closed ball $z+rB_2^n \subset \text{supp}f$. Since $\text{supp}f = \text{dom}\phi$, there is $M \in \mathbb{R}$ such that $M = \sup\{\phi(y) : y \in z + rB_2^n\}$. Thus, we have

$$f^z(x) \leq \exp\{-\sup_{y\in(z+rB_2^n)}[\langle x-z, y-z\rangle - \phi(y)]\} \leq e^M \cdot e^{-r|x-z|^2}.$$

Therefore, $\int_{\mathbb{R}^n} f^z(x)dx \leq e^M \int_{\mathbb{R}^n} e^{-r|x-z|^2}dx < \infty$. □

For any $z_1, z_2 \in \mathbb{R}^n$ and $\alpha \in (0,1)$. Let $f = e^{-\phi}$, we have $F(z) = \int_{\mathbb{R}^n} e^{-\mathcal{L}\phi(x)+\langle x,z\rangle}dx$. Since $g_x(z) := e^{-\mathcal{L}\phi(x)+\langle x,z\rangle}$ is a convex function about z, we have

$$F(\alpha z_1 + (1-\alpha)z_2) \leq \alpha F(z_1) + (1-\alpha)F(z_2). \tag{4.5}$$

If $z_1, z_2 \in \text{int dom}F$ and $z_1 \neq z_2$, then inequality (4.5) is a strict inequality. Thus $F(z)$ is strictly convex on int dom$F$.

(ii) Since $f(x)$ is even about z_0, $f(z_0+x) = f(z_0-x)$ for any $x \in \mathbb{R}^n$. For any $z \in \mathbb{R}^n$, we have

$$F(z_0+z) = \int_{\mathbb{R}^n} f^{z_0+z}(x)dx = \int_{\mathbb{R}^n} f^{z_0-z}(-x+2z_0)dx = F(z_0-z).$$

This completes the proof. □

Remark 2. *By Lemma 4.1, if f is even about z_0, then $s(f) = z_0$.*

Lemma 4.2. *Let f be a log-concave function such that $0 < \int f < \infty$, and let $G \subset \mathbb{R}^n$ be an affine subspace satisfying $G \cap \text{int supp}f \neq \emptyset$. Then there exists a unique point $z_0 \in G$ satisfying the following two equivalent claims. (i) $F(z_0) = \min\{F(z); z \in G\}$, where $F(z) := \int_{\mathbb{R}^n} f^z(x)dx$. (ii) $\text{grad}F(z_0) = \int_{\mathbb{R}^n} xf^{z_0}(x+z_0)dx \in G^\perp$.*

Proof. By Lemma 4.1, F is coercive and strictly convex on int dom$F$, thus there is a unique minimal point $z_0 = s_G(f)$ on G. Let $f = e^{-\phi}$, then $F(z) = \int_{\mathbb{R}^n} e^{-\mathcal{L}\phi(x)+\langle x,z\rangle}dx$. By the dominated convergence theorem, we have $\text{grad}F(z) = \int_{\mathbb{R}^n} xe^{-\mathcal{L}\phi(x)+\langle x,z\rangle}dx = \int_{\mathbb{R}^n} xf^z(x+z)dx$.

Next, we prove the equivalence of (i) and (ii). Let $\eta_1, \ldots, \eta_m$ $(m < n)$ be an orthonormal basis of G and let $\eta_{m+1}, \ldots, \eta_n$ be an orthonormal basis of $G^\perp$. Let $z = \sum_{i=1}^n z_i \eta_i$, since $z_0 = s_G(f) \in G$, we have $\frac{\partial F(z)}{\partial z_i}\Big|_{z=z_0} = \lim_{t\to 0} \frac{F(z_0+t\eta_i)-F(z_0)}{t} = 0, \quad i = 1, \ldots, m.$ Hence, $\mathrm{grad}F(z_0) \in G^\perp$. On the other hand, if $\mathrm{grad}F(z_0) \in G^\perp$, then $\frac{\partial F(z)}{\partial z_i}\Big|_{z=z_0} = 0,\ i = 1, \ldots, m.$ Since $F(z)$ is strictly convex on $G \cap \mathrm{int\,dom}F$, z_0 is the unique minimal point on G. □

Remark 3. *In Lemma 4.2, if $G = \mathbb{R}^n$, then the lemma shows that the Santaló point $s(f)$ of f is the barycenter of the function $f^{s(f)}$.*

Lemma 4.3. *Let f be a log-concave function such $0 < \int f < \infty$. Let $G \subset \mathbb{R}^n$ be an affine subspace satisfying $G \cap \mathrm{int\,supp}f \neq \emptyset$ and $z = s_G(f)$. Let H be an affine hyperplane such that $G \subset H$ and let g be the function defined by $g^z = S_H(f^z)$. Then we have $s_G(g) = z = s_G(f)$.*

Proof. It may be supposed that $z = s_G(f) = 0$, $H = \{(x_1, \cdots, x_n) \in \mathbb{R}^n : x_n = 0\}$ and $G = \{(x_1, \cdots, x_n) \in \mathbb{R}^n : x_{m+1} = \cdots = x_n = 0\}$ for some m, $1 \le m \le n-1$. By Lemma 4.2, we have $\int_{\mathbb{R}^n} x f^0(x)dx \in G^\perp$. Let $f^0_{x'}(t) := f^0(x' + tu)$ for any $x' \in H$, where u is the unit normal vector of H. Thus, $\int_H x_i \left(\int_{\mathbb{R}} f^0_{x'}(t)dt\right) dx' = 0$ for $1 \le i \le m$. By $g^0 = S_H(f^0)$ and (3.3), for every $x' \in H$, $\int_{\mathbb{R}} f^0_{x'}(t) = \int_{\mathbb{R}} g^0_{x'}(t)$. Thus, $\int_H x_i \left(\int_{\mathbb{R}} g^0_{x'}(t)dt\right) dx' = 0$ for $1 \le i \le m$, which conversely gives $\int_{\mathbb{R}^n} x g^0(x)dx \in G^\perp$. Thus, by Lemma 4.2 again, we obtain $s_G(g) = 0 = s_G(f)$. □

Lemma 4.4. *For a log-concave function f such that $0 < \int f < \infty$, if f is symmetric about some affine hyperplane H, then, for any $z \in H$, f^z is also symmetric about H.*

Proof. Let u be the unit normal vector of H. For any $x', y' \in H$ and $s, t \in \mathbb{R}$, since $f(x' + su) = f(x' - su)$, we have

$$f^z(y' + tu) = \inf_{x'+su\in\mathbb{R}^n} \frac{\exp\{-\langle y' + tu - z, x' + su - z\rangle\}}{f(x' + su)}$$

$$= \inf_{x'+su\in\mathbb{R}^n} \frac{\exp\{-\langle y'-z-tu, x'-z-su\rangle\}}{f(x'-su)} = f^z(y'-tu).$$

This completes the proof. □

Lemma 4.5. *Let f be a log-concave function such that $0 < \int f < \infty$ and let H be an affine hyperplane satisfying $H \cap \operatorname{int}\operatorname{supp} f \neq \emptyset$ and $z \in H \cap \operatorname{int}\operatorname{supp} f$; let λ, $0 < \lambda < 1$ such that H is λ-separating for f^z. Then*

$$\int_{\mathbb{R}^n} (S_H f)^z \geq 4\lambda(1-\lambda)\int_{\mathbb{R}^n} f^z.$$

Proof. It may be supposed that $z = 0$ and $H = \{(x_1, \ldots, x_n) : x_n = 0\}$. For $y' \in H$ and $s \in \mathbb{R}$, let (y', s) denote $y'+su$, where u is a unit normal vector of H. For f^0 and $s \in \mathbb{R}$, we define a new function

$$f^0_{(s)}(y') := f^0(y', s), \text{ for any } y' \in H.$$

Next we shall prove that for any $y' \in H$ and $s, t > 0$

$$\left(\frac{t}{s+t}\cdot f^0_{(s)}\right) \star \left(\frac{s}{s+t}\cdot f^0_{(-t)}\right)(y') \leq (S_H f)^0_{(\frac{2st}{s+t})}(y'). \tag{4.6}$$

Claim 2. For any $x' \in H$ and $w \in \mathbb{R}$, if $(S_H f)(x'+wu) > 0$, then there is some $w_1 \in \mathbb{R}$ such that $(S_H f)(x'+wu) \leq f(x'+w_1 u)$ and $(S_H f)(x'+wu) \leq f(x'+(w_1-2w)u)$.

Proof of Claim 2. Let $f = e^{-\phi}$, since $(S_H f)(x'+wu) > 0$, then $(S_H\phi)(x'+wu) < +\infty$. By Proposition 1(iii), there is $w_1 \in \mathbb{R}$ such that $(S_H\phi)(x'+wu) \geq \phi(x'+w_1 u)$ and $(S_H\phi)(x'+wu) \geq \phi(x'+(w_1-2w)u)$, here we assume $\phi(x'+w_1 u)$ or $\phi(x'+(w_1-2w)u)$ equals the limit in Proposition 1(iii), which doesn't affect our proof. Hence the claim follows. □

For any y'_1, $y'_2 \in H$ such that $y' = y'_1 + y'_2$, we have

$$\begin{aligned}(S_H f)^0_{(\frac{2st}{s+t})}(y') &= \inf_{(x',w)\in H\times\mathbb{R}} \frac{\exp\{-\langle (y', \frac{2st}{s+t}), (x', w)\rangle\}}{(S_H f)(x', w)} \\ &\geq \inf_{(x',w)\in H\times\mathbb{R}} \frac{\exp\{-\langle (y', \frac{2st}{s+t}), (x', w)\rangle\}}{f(x', w_1)^{\frac{t}{s+t}} f(x', w_1-2w)^{\frac{s}{s+t}}}\end{aligned}$$

$$\begin{aligned}
&\geq \inf_{(x',w)\in H\times\mathbb{R}} \frac{\exp\{-\frac{t}{s+t}\langle(\frac{s+t}{t}y_1', s), (x', w_1)\rangle\}}{f(x', w_1)^{\frac{t}{s+t}}} \\
&\quad \times \inf_{(x',w)\in H\times\mathbb{R}} \frac{\exp\{-\frac{s}{s+t}\langle(\frac{s+t}{s}y_2', -t), (x', w_1-2w)\rangle\}}{f(x', w_1-2w)^{\frac{s}{s+t}}} \\
&\geq f^0\left(\frac{s+t}{t}y_1', s\right)^{\frac{t}{s+t}} f^0\left(\frac{s+t}{s}y_2', -t\right)^{\frac{s}{s+t}},
\end{aligned}$$

where the first inequality is by Claim 2, and the second inequality is by $\inf(AB) \geq (\inf A)(\inf B)$, and last inequality is by the definition of the polar of functions. Since y_1' and y_2' are arbitrary, we get (4.6).

Let $F_0(w) = \int_H (S_H f)^0_{(w)}$, $F_1(s) = \int_H f^0_{(s)}$ and $F_2(t) = \int_H f^0_{(-t)}$. By the Prékopa inequality and (4.6), we have

$$F_0(\frac{2st}{s+t}) \geq F_1(s)^{\frac{t}{s+t}} F_2(t)^{\frac{s}{s+t}} \text{ for every } s, t > 0.$$

Now, by Proposition 1(i) and Lemma 4.4, $(S_H f)^0$ is symmetric about H, we have $\int_0^{+\infty} F_0 = \frac{1}{2}\int_{\mathbb{R}^n} (S_H f)^0$ and since H is λ-separating for f^0, we have $\left(\int_0^{+\infty} F_1\right)\left(\int_0^{+\infty} F_2\right) = \lambda(1-\lambda)\left(\int_{\mathbb{R}^n} f^0\right)^2$. Since F_0, F_1, $F_2 : [0, +\infty) \to \mathbb{R}^+$ satisfy the hypothesis of Lemma 2.1, and by definitions of F_1 and F_2, one has $\int_0^{+\infty} F_1 + \int_0^{+\infty} F_2 = \int_{\mathbb{R}^n} f^0$, thus, by Lemma 2.1

$$\frac{2}{\int_{\mathbb{R}^n}(S_H f)^0} \leq \frac{1}{2}\left(\frac{1}{\int_0^{+\infty} F_1} + \frac{1}{\int_0^{+\infty} F_2}\right) = \frac{1}{2\lambda(1-\lambda)\int_{\mathbb{R}^n} f^0}.$$

This gives the desired inequality. □

Lemma 4.6. *If f is an integrable, unconditional, log-concave function, then $\int_{\mathbb{R}^n} f \int_{\mathbb{R}^n} f^0 \leq (2\pi)^n$.*

Proof. Let $f_1 = f$, $f_2 = f^0$ and $f_3 = e^{-\frac{|x|^2}{2}}$, then f_1, f_2 and f_3 are unconditional. Thus we have $\int_{\mathbb{R}^n} f_j = 2^n \int_{\mathbb{R}^n_+} f_j$, $j = 1, 2, 3$. For $(y_1, \ldots, y_n) \in \mathbb{R}^n$, we define $g_i(y_1, \ldots, y_n) = f_i(e^{y_1}, \ldots, e^{y_n}) e^{\sum_{i=1}^n y_i}$. We get $\int_{\mathbb{R}^n_+} f_j = \int_{\mathbb{R}^n} g_j$, and for every $s, t \in \mathbb{R}^n$, $g_1(s)g_2(t) \leq g_3\left(\frac{s+t}{2}\right)^2$. Hence $\int_{\mathbb{R}^n} f \int_{\mathbb{R}^n} f^0 \leq (2\pi)^n$ follows from Prékopa inequality. □

Proof of Theorem 1.2. We proceed by n successive Steiner symmetrizations until we get an unconditional log-concave function.

Let $u_1 \in S^{n-1}$, u_1 orthogonal to $H = H_1$ and let $(u_i)_{i=2}^n \subset S^{n-1}$ such that $(u_1, \dots, u_n)$ form an orthonormal basis for $\mathbb{R}^n$. Let $z_1 = s_{H_1}(f)$ and define a log-concave function f_1 by the identity $f_1^{z_1} = S_{H_1}(f^{z_1})$. Then $\int f_1^{z_1} = \int f^{z_1}$. By Proposition 1(i) and Lemma 4.4, f_1 is symmetric about H_1 and by Lemma 4.5, applied to f^{z_1}, $z = z_1$ and $H = H_1$, λ-separating for $f = (f^{z_1})^{z_1}$, we get $\int_{\mathbb{R}^n} f_1 \geq 4\lambda(1-\lambda)\int_{\mathbb{R}^n} f$ and thus $\int f_1 \int f_1^{z_1} \geq 4\lambda(1-\lambda)\int f \int f^{z_1}$. Choose now the hyperplane H_2, orthogonal to u_2, and medial for f_1 and define $z_2 = s_{(H_1\cap H_2)}(f_1)$. By Lemma 4.3 we have $z_1 = s_{H_1}(f) = s_{H_1}(f_1)$, we get $\int f_1^{z_2} = \min_{z\in H_1\cap H_2} \int f_1^z \geq \min_{z\in H_1} \int f_1^z = \int f_1^{z_1}$. We define now a new log-concave function f_2 by the identity $f_2^{z_2} = S_{H_2}(f_1^{z_2})$. By Proposition 1(ii) and Lemma 4.4, f_2 is symmetric about both H_1 and H_2. Since H_2 is medial for f_1, we get by Lemma 4.5 applied to $f_1^{z_2}$, $z = z_2$ and $H = H_2$ that $\int f_2 \geq \int f_1$. Moreover, we have $\int f_2^{z_2} = \int S_{H_2}(f_1^{z_2}) = \int f_1^{z_2} \geq \int f_1^{z_1}$. It follows that $\int f_2 \int f_2^{z_2} \geq \int f_1 \int f_1^{z_1}$.

We continue this procedure by choosing hyperplanes $H_2, \dots, H_n$, points $z_2, \dots, z_n$, and defining log-concave functions $f_2, \dots, f_n$ such that for $2 \leq i \leq n$, we have (i) H_i is medial for f_{i-1} and orthogonal to u_i; (ii) $z_i = s_{(H_1\cap H_2\cap\cdots\cap H_i)}(f_{i-1})$; (iii) $f_i^{z_i} = S_{H_i}(f_{i-1}^{z_i})$. From (ii) (iii) and Lemma 4.3, we have $z_i = s_{(H_1\cap\cdots\cap H_i)}(f_{i-1}) = s_{(H_1\cap\cdots\cap H_i)}(f_i)$. Choosing H_{i+1}, z_{i+1}, f_{i+1} according to (i) (ii) (iii), we get thus $\int f_{i+1}^{z_{i+1}} = \int S_{H_{i+1}}(f_i^{z_{i+1}}) = \int f_i^{z_{i+1}} \geq \int f_i^{s_{(H_1\cap\cdots\cap H_i)}(f_i)} = \int f_i^{z_i}$. Now, Lemma 4.5 applied to $f_i^{z_{i+1}}$, $z = z_{i+1}$ and H_{i+1}, medial for $f_i = (f_i^{z_{i+1}})^{z_{i+1}}$, gives $\int f_{i+1} \geq \int f_i$. Thus, $\int f_i \int f_i^{z_i}$ is an increasing sequence, for $2 \leq i \leq n$. Therefore, we have $4\lambda(1-\lambda)\int f \int f^{z_1} \leq \int f_1 \int f_1^{z_1} \leq \cdots \leq \int f_n \int f_n^{z_n}$. From Proposition 1(ii), f_n is an unconditional function about z_n and $z_n \in H_1 \cap H_2 \cap \cdots \cap H_n$ is a center of symmetry for f_n. By Lemma 4.6, we have $\int f \int f^{z_1} \leq \frac{(2\pi)^n}{4\lambda(1-\lambda)}$, this concludes the proof. $\square$

References

[1] S. Artstein, B. Klartag, V. D. Milman, *On the Santaló point of a function and a functional Santaló inequality*, Mathematika 54 (2004), 33-48.

[2] K. Ball, *Isometric problems in l_p and sections of convex sets*, Doctoral thesis, University of Cambridge, 1986.

[3] K. Ball, *Logarithmically concave functions and sections of convex sets in $\mathbb{R}^n$*, Studia Math. 88 (1988), 69-84.

[4] W. Blaschke, *über affine Geometrie 7: Neue Extremeigenschaften von Ellipse und Ellipsoid.* Wilhelm Blaschke Gesammelte Werke 3. Thales Verlag, Essen (1985)

[5] H. J. Brascamp, E. H. Lieb, J. M. Luttinger, *A General Rearrangement Inequality for Multiple Integrals*, J. Funct. Anal. 17 (1974), 227-237.

[6] A. Burchard, *Steiner symmetrization is continuous in $W^{1,p}$*, Geom. Funct. Anal. 7 (1997), 823-860.

[7] A. Burchard, *A short course on rearrangement inequalities*, available at http://www.math.utoronto.ca/almut/rearrange.pdf, 2009.

[8] M. Fortier, *Convergence results for rearrangements: Old and new*, M.S. Thesis, University of Toronto, December 2010.

[9] M. Fradelizi, M. Meyer, *Some functional forms of Blaschke-Santaló inequality*, Math. Z. 256 (2007), 379-395.

[10] J. Lehec, *The symmetric property τ for the Gaussian measure*, Ann. Fac. Sci. Toulouse Math. 17(6) (2008), 357-370.

[11] J. Lehec, *Partitions and functional Santaló inequality*, Arch. Math. 92 (2009), 89-94.

[12] J. Lehec, *A direct proof of the functional Santaló inequality*, C. R. Math. Acad. Sci. Paris, 347 (2009), 55-58.

[13] E. Lutwak, G. Zhang, *Blaschke-Santaló inequalities*, J. Differ. Geom. 47(1) (1997), 1-16.

[14] E. Lutwak, D. Yang, G. Zhang, *Moment-entropy inequalities*, Ann. Probab. 32 (2004), 757-774.

[15] M. Meyer, A. Pajor, *On the Blaschke Santaló inequality*, Arch. Math. 55 (1990), 82-93.

[16] A. Prkopa, *Logarithmic concave measures with applications to stochastic programming*, Acta Sci. Math. (Szeged) 32 (1971), 301-316.

[17] A. Prkopa, *On logarithmic concave measures and functions*, Acta Sci. Math. (Szeged) 34 (1973), 339-343.

[18] L. A. Santaló, *An affine invariant for convex bodies of n-dimensional space*. Port. Math. 8 (1949), 155-161.

[19] R. Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, Encyclopedia Math. Appl., vol. 44, Cambridge University Press, Cambridge, 1993.

School of Mathematical Sciences, Peking University, Beijing, 100871, China; Department of Mathematics, Department of Mathematics, Shanghai University, Shanghai, 200444, China

E-mail address: lxyoujiang@126.com

Department of Mathematics, Shanghai University, Shanghai, 200444, China

E-mail address: gleng@staff.shu.edu.cn